

THIS REPORT HAS BEEN DELIMITED
AND CLEARED FOR PUBLIC RELEASE
UNDER DOD DIRECTIVE 5200.20 AND
NO RESTRICTIONS ARE IMPOSED UPON
ITS USE AND DISCLOSURE.

DISTRIBUTION STATEMENT A

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED.

AD 81483

Armed Services Technical Information Agency

**Reproduced by
DOCUMENT SERVICE CENTER
KNOTT BUILDING, DAYTON, 2, OHIO**

This document is the property of the United States Government. It is furnished for the duration of the contract and shall be returned when no longer required, or upon recall by ASTIA to the following address:
Armed Services Technical Information Agency, Document Service Center,
Knott Building, Dayton 2, Ohio.

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

UNCLASSIFIED

AD No 81483


ASTIA FILE COPY

FC

REPORT NO. HE-150-133

TECHNICAL REPORT

UNIVERSITY OF CALIFORNIA
INSTITUTE OF ENGINEERING RESEARCH
BERKELEY, CALIFORNIA



STUDIES OF BOUNDARY LAYER SLIP SOLUTIONS AND ALDEN'S
METHOD FOR BOUNDARY LAYER CORRECTION

by

S. BELL

SERIES NO. 20.....

ISSUE NO. 106.....

DATE December 6, 1955.....

CONTRACT N7-onr-295-Task 3
PROJECT NR 061-003
REPORT NO. HE-150-133
SERIES NO. 20-106
DECEMBER 6, 1955

JOINTLY SPONSORED BY
OFFICE OF NAVAL RESEARCH AND
OFFICE OF SCIENTIFIC RESEARCH

Reproduction in whole or in part
is permitted for any purpose of
the United States Government

FLUID FLOW AND
HEAT TRANSFER
AT LOW PRESSURES
AND TEMPERATURE

STUDIES OF BOUNDARY LAYER SLIP SOLUTIONS AND ALDEN'S
METHOD FOR BOUNDARY LAYER CORRECTION

by

S. BELL

FACULTY INVESTIGATORS:

S. A. SCHAAF, ASSOCIATE PROFESSOR OF ENGINEERING SCIENCE

L. TALBOT, ASSISTANT PROFESSOR OF MECHANICAL ENGINEERING

Approved by:

L. Talbot

Note: This report was condensed from a Ph.D. thesis, University of California,
Berkeley, 1955.

TABLE OF CONTENTS

	Page
Abstract	iii
Nomenclature	iv
1.0 Introduction	1
2.0 Analysis	2
3.0 Conclusions	15
Appendix I	16
Appendix II	17
Appendix III	21
List of References	23

ABSTRACT

Two aspects of incompressible laminar boundary layer flow on a semi-infinite flat plate at zero incidence are discussed. Firstly, Alden's proposed scheme of solution of the Navier-Stokes equations for this problem in terms of an expansion in inverse powers of the local Reynolds number is demonstrated to be invalid, inasmuch as the boundary conditions in the free stream cannot be satisfied to all orders. Secondly, an account is given of the effect of a velocity slip boundary condition on the solution of the Oseen boundary layer equations for the same problem. It is found that an additional non-uniformity in the solution is caused by the introduction of slip.

NOMENCLATURE

C_D	Drag coefficient, eq. (44)
C_f	Local skin friction coefficient, eq. (42)
C_{11}, C_{22}	Constants defined in eq. (18)
C_{ni}	Constants defined in eqs. (23) and (30)
$F_n(\xi)$	Functions defined in eq. (14)
$F_{ni}(\xi)$	Functions defined in eqs. (24), (25), (28) and (30)
$f_n(\xi)$	Functions defined in eq. (12)
$g_n(\xi)$	Functions defined in eq. (58)
$h_n(\xi)$	Functions defined in eq. (26)
K	Constant defined in eq. (32) ($K = K_{23}$)
K_n	Knudsen number
K_{nn}	Constants of integration (Appendix II)
$O(x^{-N})$	Symbolizes terms of order $-N$, i.e., $\lim_{x \rightarrow \infty} x^N O(x^{-N}) = \text{constant}$
p	Pressure
r	$= \sqrt{x^2 + y^2}$
Re_L	$= \frac{u_\infty L}{\nu_\infty}$, Reynolds number based on L
Re_x	$= \frac{u_\infty x}{\nu_\infty}$, Reynolds number based on x
s	$= u_\infty / \sqrt{2RT_\infty}$, molecular speed ratio
u, v	x and y velocity components
$u_n(x, y)$	Functions defined in eq. (46)
u_∞	Free stream velocity
x, y	Cartesian coordinates
α_n	Constants defined in eq. (29)
β	Constant defined in eq. (21)
Γ	Gamma function

NOMENCLATURE (continued)

ζ	Slip parameter, see eq. (36)
$\eta = \tau y^{-\frac{1}{2}}$, auxiliary variable
$\theta = \tan^{-1} y/x$	
μ	Viscosity
$\nu = \mu/\rho$, kinematic viscosity
ν_∞	Free stream kinematic viscosity
$\xi = \eta u_\infty^{-\frac{1}{2}}$, auxiliary variable
ρ	Density
ρ_∞	Free stream density
σ, τ	Parabolic coordinates, eq. (6)
ψ	Stream function, eq. (2)
$\psi_n(\sigma, \eta)$	Functions defined in eq. (11)
∇^2	Laplacian operator
\sim	Asymptotic equality

1.0 INTRODUCTION

Blasius solved the incompressible boundary layer problem for laminar flow past a semi-infinite flat plate at zero angle of attack (references 1 and 2) and his result describes the flow quite well for regions where the local Reynolds number, $u_{\infty} x / \nu$, is large. Von Karman has proposed (reference 3) that the Blasius solution be considered the first term of an asymptotic representation of a solution to the full Navier-Stokes equations; calculation of the second term should extend boundary layer theory to lower Reynolds numbers. Due to the non-uniformities which are inherent in boundary layer theory, any such representation will contain terms which express these non-uniformities; in particular, one would not expect a simple power series in the viscosity. The specific form for the stream function expansion suggested by von Karman has a line singularity on the y-axis (the plate is assumed to occupy the positive x-axis with the leading edge at the origin). Alden (reference 4) improved this situation somewhat by modifying von Karman's representation in such a way as to rotate the line singularity to the negative x-axis. Adopting von Karman's philosophy, he obtained a "first correction" to the Blasius solution. At first glance, Alden's iteration scheme seems to lead to an asymptotic series which is formally a solution of the Navier-Stokes problem. In the first part of this paper it will be shown that Alden's method cannot lead to such a solution because of an unavoidable violation of the free stream boundary conditions.

In the case of a gas at low Reynolds numbers the boundary conditions at a solid surface must be modified to account for the slipping of the gas at the wall (references 5 through 9). To support the conjecture that slip introduces an additional non-uniformity into the solution, the linearized boundary layer problem (for the flat plate) with a slip condition is examined in detail in the second part of this paper. An explicit solution is known and this is compared with the divergent series solution obtained by a perturbation in the "slip parameter".

2.0 ANALYSIS

2.1 Alden's Method

The Navier-Stokes equations of motion and continuity, specialized for the steady state two dimensional flow of an incompressible fluid are

$$\left. \begin{aligned} u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} &= -\frac{1}{\rho} \frac{\partial p}{\partial x} + \nu \nabla^2 u \\ u \frac{\partial v}{\partial x} + v \frac{\partial v}{\partial y} &= -\frac{1}{\rho} \frac{\partial p}{\partial y} + \nu \nabla^2 v \\ \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} &= 0 \end{aligned} \right\} \quad (1)$$

where x and y are cartesian coordinates, u and v are the cartesian velocity components, ∇^2 is the Laplacian operator, and ρ , p and ν are, respectively, the density, the pressure and the kinematic viscosity.

A stream function, ψ , may be introduced so that

$$u = \frac{\partial \psi}{\partial y} ; \quad v = -\frac{\partial \psi}{\partial x} \quad (2)$$

and the continuity equation is automatically satisfied. If the momentum equations are cross differentiated and subtracted, to eliminate the pressure terms, one obtains, in terms of the stream function

$$\frac{\partial \psi}{\partial y} \frac{\partial}{\partial x} \nabla^2 \psi - \frac{\partial \psi}{\partial x} \frac{\partial}{\partial y} \nabla^2 \psi = \nu \nabla^4 \psi \quad (3)$$

Physical arguments lead one to suspect that the boundary layer equation (reference 10)

$$\frac{\partial \psi}{\partial y} \frac{\partial^3 \psi}{\partial x \partial y^2} - \frac{\partial \psi}{\partial x} \frac{\partial^3 \psi}{\partial y^3} = \nu \frac{\partial^4 \psi}{\partial y^4} \quad (4)$$

may describe the flow quite well in regions where the local Reynolds number, $u_\infty x / \nu$, is large, (u_∞ being the free stream velocity).

To complete this stream function formulation of the incompressible boundary layer problem for laminar flow past a semi-infinite flat plate at zero angle of attack, it will be assumed that the plate occupies the positive x-axis with the leading edge at the origin and it will be required that

$$\left. \begin{aligned} \psi = \frac{\partial \psi}{\partial x} = \frac{\partial \psi}{\partial y} = 0 \quad \text{at } y = 0, x > 0 \\ \frac{\partial \psi}{\partial x} \rightarrow 0 \\ \frac{\partial \psi}{\partial y} \rightarrow u_{\infty} \end{aligned} \right\} \text{ as } r \rightarrow \infty, 0 < \theta < \pi \quad (5)$$

where r and θ are polar coordinates with origin at the leading edge of the plate. This is essentially the problem solved by Blasius.

In terms of parabolic coordinates,

$$x + iy = (\sigma + i\tau)^2 = r e^{i\theta} \quad (6)$$

equation (3) becomes

$$\begin{aligned} & (\sigma^2 + \tau^2) \left\{ \frac{\partial \psi}{\partial \tau} \frac{\partial}{\partial \sigma} \nabla^2 \psi - \frac{\partial \psi}{\partial \sigma} \frac{\partial}{\partial \tau} \nabla^2 \psi \right\} \\ & - 2 \nabla^2 \psi \left\{ \sigma \frac{\partial \psi}{\partial \tau} - \tau \frac{\partial \psi}{\partial \sigma} \right\} \\ & = \nu \left\{ 4 - 4 \left(\sigma \frac{\partial}{\partial \sigma} + \tau \frac{\partial}{\partial \tau} \right) + (\sigma^2 + \tau^2) \nabla^2 \right\} \nabla^2 \psi \end{aligned} \quad (7)$$

$$\text{where } \nabla^2 = \frac{\partial^2}{\partial \sigma^2} + \frac{\partial^2}{\partial \tau^2}$$




Figure 1.

Applying the analogous boundary layer arguments to equation (7) one obtains the "parabolic boundary layer equation",

$$\sigma^2 \frac{\partial \psi}{\partial \tau} \frac{\partial^3 \psi}{\partial \sigma \partial \tau^2} - \sigma^2 \frac{\partial \psi}{\partial \sigma} \frac{\partial^3 \psi}{\partial \tau^3} - 2\sigma \frac{\partial^2 \psi}{\partial \tau^2} \frac{\partial \psi}{\partial \tau} = \nu \sigma^2 \frac{\partial^4 \psi}{\partial \tau^4} \quad (8)$$

Note that the parabolic boundary layer equation is not the same as the cartesian boundary layer equation expressed in parabolic coordinates (see reference 11).

H. L. Alden (reference 4), following a suggestion by von Karman (reference 3), attempted to improve the boundary layer solution by developing the iteration scheme which solves equation (1) with the Blasius solution as the first term in an expansion. Using his scheme, Alden found two terms in a proposed expansion. The first term is the solution of the parabolic boundary layer problem and the second term may be thought of as a correction term. Since there is no obvious mathematical reason for concluding that his scheme will not lead to an asymptotic expansion which is formally a solution of the problem, it is worth while to investigate further. Alden introduces the auxiliary variables

$$\left. \begin{aligned} \eta &= \tau \nu^{-\frac{1}{2}} \\ \xi &= \eta u_\infty^{-\frac{1}{2}} \end{aligned} \right\} \quad (9)$$

and rewrites equation (7) as

$$\begin{aligned}
 & (\sigma^2 + \nu\eta^2) \left\{ \frac{\partial \psi}{\partial \eta} \left(\frac{\partial^3 \psi}{\partial \sigma^3} + \nu^{-1} \frac{\partial^3 \psi}{\partial \sigma \partial \eta^2} \right) - \frac{\partial \psi}{\partial \sigma} \left(\frac{\partial^3 \psi}{\partial \sigma^2 \partial \eta} + \nu^{-1} \frac{\partial^3 \psi}{\partial \eta^3} \right) \right\} \\
 & - 2 \left(\frac{\partial^2 \psi}{\partial \sigma^2} + \nu^{-1} \frac{\partial^2 \psi}{\partial \eta^2} \right) \left(\sigma \frac{\partial \psi}{\partial \eta} - \nu \eta \frac{\partial \psi}{\partial \sigma} \right) \\
 & - \nu^{\frac{1}{2}} \left\{ 4 \left(\frac{\partial^2 \psi}{\partial \sigma^2} + \nu^{-1} \frac{\partial^2 \psi}{\partial \eta^2} \right) - 4 \sigma \left(\frac{\partial^3 \psi}{\partial \sigma^3} + \nu^{-1} \frac{\partial^3 \psi}{\partial \sigma \partial \eta^2} \right) \right. \\
 & \quad \left. - 4 \eta \left(\frac{\partial^3 \psi}{\partial \sigma^2 \partial \eta} + \nu^{-1} \frac{\partial^3 \psi}{\partial \eta^3} \right) \right. \\
 & \quad \left. + (\sigma^2 + \nu\eta^2) \left(\frac{\partial^4 \psi}{\partial \sigma^4} + 2\nu^{-1} \frac{\partial^4 \psi}{\partial \sigma^2 \partial \eta^2} + \nu^{-2} \frac{\partial^4 \psi}{\partial \eta^4} \right) \right\} \\
 & = 0
 \end{aligned} \tag{10}$$

Alden proposed the iteration scheme whereby $\psi(\sigma, \eta, \nu)$ is formally expressed as

$$\psi(\sigma, \eta, \nu) = \nu^{\frac{1}{2}} \psi_1(\sigma, \eta) + \nu^{\frac{3}{2}} \psi_2(\sigma, \eta) + \nu^{\frac{5}{2}} \psi_3(\sigma, \eta) + \dots \tag{11}$$

and this is to be substituted into equation (10) and the coefficients of explicit powers of ν set equal to zero. The function, $\nu^{\frac{1}{2}} \psi_1(\sigma, \eta)$, will be found to be a solution of the parabolic boundary layer problem and this scheme may be thought of as a perturbation, in ν , around the parabolic boundary layer solution. If, in equation (7), ν is set equal to zero, the order of the equation is depressed. One therefore calls this perturbation scheme singular and expects some sort of singularity in the solution (references 12 and 13). Furthermore, the expansion (11) is not a true power series in ν since the coefficients $\psi_n(\sigma, \eta)$ depend implicitly on ν through η .

Alden's scheme produces a system of equations for the $\psi_n(\sigma, \eta)$ and for each equation the variables may be separated if one sets

$$\psi_n(\sigma, \eta) = (u_\infty \sigma^2)^{\frac{3-2n}{2}} f_n(\xi) \quad (12)$$

This yields a system of ordinary differential equations for the $f_n(\xi)$. Unfortunately expression (12) implies that, for $n \geq 2$,

$\psi_n(\sigma, \eta) \rightarrow \infty$ as $\sigma \rightarrow 0$. It might be possible however, to show that the expression (11) is an asymptotic series, uniform in σ and η , in some region excluding an arbitrarily small region bounded by a parabola enclosing the stagnation line ($\sigma = 0$) and the leading edge. This introduces the further complication of whether the boundary conditions which remain are sufficient for the problem. The region mentioned above excludes the origin, which is the most likely point where a non-uniformity will occur in the solution.

The iteration scheme yields the following equations for the $f_n(\xi)$

$$(f_1''' + f_1 f_1'')' = 0 \quad (13)$$

and for $n = 2, 3, 4, \dots$

$$f_n''' + f_1 f_n'' + (2n-1)f_1' f_n'' + f_1'' f_n' + (3-2n)f_1''' f_n = F_n(\xi) \quad (14)$$

where $F_n(\xi)$ is dependent on

ξ, f_1, f_2, \dots and f_{n-1} ,

and the primes indicate differentiation with respect to ξ .

($F_2(\xi), F_3(\xi)$, and $F_4(\xi)$ are exhibited in Appendix I).

The boundary conditions (5) require the following behavior for the $f_n(\xi)$, introduced in equations (11) and (12),

$$\left. \begin{aligned} f_n(0) &= 0 \\ f_n'(0) &= 0 \end{aligned} \right\} n = 1, 2, 3, \dots \quad (16)$$

$$\left. \begin{aligned} f_1(\xi)/2\xi &\rightarrow 1 \\ f_1'(\xi)/2 &\rightarrow 1 \\ f_n(\xi)/\xi &\rightarrow 0 \\ f_n'(\xi) &\rightarrow 0 \end{aligned} \right\} \begin{matrix} n = 2, 3, 4, \dots \\ \text{as } \xi \rightarrow \infty \end{matrix} \quad (16)$$

From equation (13)

$$f_1''' + f_1 f_1'' = C_{11}, \text{ a constant.} \quad (17)$$

Alden states that because of conditions (16)

$$f_1'''(\xi) \rightarrow 0 \text{ and } f_1(\xi)f_1''(\xi) \rightarrow 0 \text{ as } \xi \rightarrow \infty,$$

This can be proved by making the highly reasonable assumption that the velocity gradients are continuous. Because of this hypothesis, $f_1(\xi)$ and its first two derivatives are continuous on $0 \leq \xi < \infty$. Now, rewrite equation (17) as follows.

$$\frac{f_1''(\xi)}{\xi} + \frac{f_1(\xi)f_1'(\xi)}{\xi} = \frac{C_{12}}{\xi} + C_{11} + \frac{1}{\xi} \int_0^\xi [f_1'(\eta)]^2 d\eta \quad (18)$$

Using conditions (16), it is easy to show that

$$\lim_{\xi \rightarrow \infty} \frac{f_1''(\xi)}{\xi} = C_{11} \quad (19)$$

$$\lim_{\xi \rightarrow \infty} \frac{f_1'(\xi)}{\xi^2} = C_{11} = 0$$

Thus, the equation for f_1 becomes

$$f_1''' + f_1 f_1'' = 0 \quad (20)$$

This equation with the conditions in equations (16) possesses a solution (reference 14), often referred to as the Blasius function, which is, together with all of its derivatives, continuous. Furthermore, this is the only solution with continuous derivatives (reference 14).

The Blasius function exhibits the following asymptotic behavior as $\xi \rightarrow \infty$ (reference 15).

$$\begin{aligned}
 f_1(\xi) &\sim 2\xi - \beta, & \beta &= 1.73 \\
 f_1'(\xi) &\sim 2 \\
 f_1^{(n)}(\xi) &\sim 0, & n &= 2, 3, 4, \dots
 \end{aligned}
 \tag{21}$$

Using equations (6), (9) and (12), one may formally rewrite expression (11):

$$\psi(\sigma, \eta, \nu) = \frac{u_\infty y}{2\xi} \sum_{n=0}^{\infty} f_{n+1}(\xi) \left(\frac{\nu}{u_\infty \sigma^2} \right)^n \tag{22}$$

If conditions (16) are satisfied and each $f_n(\xi)$ is continuous on $0 \leq \xi < \infty$, then $u_\infty y f_n(\xi) / 2\xi$ is bounded on $0 \leq \xi < \infty$ and expression (22) is an asymptotic expansion.

Unfortunately, Alden's scheme does not lead to a solution of the Navier-Stokes problem since $f_4(\xi)$ cannot be made to satisfy conditions (16). In particular, it is impossible to choose $f_4(\xi)$ so that $\lim_{\xi \rightarrow \infty} \frac{f_4(\xi)}{\xi} = 0$. This will be shown in the following pages.

$$\begin{aligned}
 \text{Let } C_{n1} &= f_n'''(0) \\
 C_{n2} &= f_n''(0)
 \end{aligned}
 \tag{23}$$

One may rewrite equation (14) as

$$f_n''' + f_1 f_n'' + (2n-2)f_1' f_n' + (3-2n)f_1'' f_n = F_{n1}(\xi) \tag{24}$$

$$F_{n1}(\xi) = C_{n1} + \int_0^\xi F_n(\eta) d\eta$$

Integrating again,

$$f_n'' + f_1 f_n' + (2n-3)f_1' f_n = F_{n2}(\xi) \tag{25}$$

$$F_{n2}(\xi) = C_{n2} + \int_0^\xi [F_{n1}(\eta) + 2(2n-3)f_1''(\eta)f_n(\eta)] d\eta$$

Let

$$h_n(\xi) = \exp\left[(\xi - \beta/2)^2\right] \frac{d^{2n-4}}{d\xi^{2n-4}} \left[-(\xi - \beta/2)^2\right] \tag{26}$$

for $n = 2, 3, 4, \dots$

Multiply equation (25) by $h_n(\xi)$ and obtain

$$[h_n f_n' + (h_n f_1 - h_n') f_n]' + f_n [h_n'' - f_1 h_n' + 2(n-2) f_1' h_n] = h_n F_{n2} \quad (27)$$

One can show that

$$h_n f_n' + (h_n f_1 - h_n') f_n = F_{n3}(\xi), \quad (28)$$

$$F_{n3}(\xi) = \int_0^\xi [h_n(\eta) F_{n2}(\eta) + f_n(\eta) \{ (f_1(\eta) - 2\eta + \beta) h_n'(\eta) + 2(n-2)(2 - f_1'(\eta)) h_n(\eta) \}] d\eta,$$

and that α_n may be chosen so, that, for $\alpha_n \leq \xi < \infty$,

$$\begin{aligned} \left[\frac{e^{(\xi - \beta/2)^2}}{h_n} f_n \right]' &= \frac{e^{(\xi - \beta/2)^2}}{h_n^2} [h_n f_n' + (2\xi - \beta) h_n f_n - h_n' f_n] \\ &= \frac{e^{(\xi - \beta/2)^2}}{h_n^2} [F_{n3}(\xi) + (2\xi - \beta - f_1) h_n f_n] \end{aligned} \quad (29)$$

$$\begin{aligned} f_n(\xi) &= C_{n4} h_n(\xi) e^{-(\xi - \beta/2)^2} \\ &\quad + h_n(\xi) e^{-(\xi - \beta/2)^2} \int_{\alpha_n}^\xi e^{(\eta - \beta/2)^2} F_{n4}(\eta) d\eta, \end{aligned} \quad (30)$$

$$F_{n4}(\xi) = \frac{1}{[h_n(\xi)]^2} \left[F_{n3}(\xi) + h_n(\xi) f_n(\xi) (2\xi - \beta - f_1(\xi)) \right]$$

The $h_n(\xi)$ and $f_n(\xi)$ are asymptotically finite, i.e., for each n there is an integer q such that $h_n(\xi)/\xi^q \rightarrow 0$ as $\xi \rightarrow \infty$.

$$\begin{aligned}
f_n(\xi) &\sim h_n(\xi) e^{-(\xi-\beta/2)^2} \int_{\eta_n}^{\xi} e^{(\eta-\beta/2)^2} F_{n+1}(\eta) d\eta \\
F_{n+1}(\xi) &\sim [h_n(\xi)]^{-2} F_n(\xi) \\
F_n(\xi) &\sim \int_0^{\xi} h_n(\eta) F_{n-1}(\eta) d\eta + \text{a constant} \\
F_{n-1}(\xi) &\sim C_{n-1} + \int_0^{\xi} F_{n-2}(\eta) d\eta + \text{a constant} \\
F_{n-2}(\xi) &\sim C_{n-2} + \int_0^{\xi} F_{n-3}(\eta) d\eta
\end{aligned} \tag{31}$$

Carrying out the obvious algebra, one can show (see Appendix II) that

$$f_2(\xi) \sim K e^{-(\xi-\beta/2)^2} \int_0^{\xi} e^{(\eta-\beta/2)^2} d\eta \tag{32}$$

and

$$\lim_{\xi \rightarrow \infty} \frac{f_4(\xi)}{\xi^2 \ln \xi} = 6\beta K \tag{33}$$

Expression (32) is identical with Alden's result and is introduced here to show that $K \neq 0$. In fact Alden computed $K = 1.660$ by numerical integration. Thus, $f_4(\xi)$ cannot be made to satisfy condition (16) and consequently the proposed expressions (11) and (12) cannot be made, even formally, to satisfy equations (10) and (5). It may be remarked that the Alden approach with a slip boundary condition is subject to the same difficulty.

2.2 The Flat Plate Boundary Layer Problem With Slip

In an attempt to incorporate rarefaction effects into boundary layer theory, one is led (reference 9) to consider a slip boundary condition,

$$u = \xi \frac{\partial u}{\partial y}, \text{ where } \xi \text{ will be called the slip parameter. } \xi \text{ is}$$

proportional to the mean free path length.

The flat plate boundary layer problem with slip (i.e., with a slip boundary condition) has not yet been solved. With the idea of gaining some knowledge of the slip effect in general, the corresponding linearized problem for incompressible flow, which has been solved explicitly (reference 7), will now be examined. First, the problem will be formulated and the solution displayed. Secondly, a series solution of restricted validity will be obtained by a perturbation scheme in ζ . Lastly, these solutions will be compared and analyzed.

The linearized, incompressible, steady-state boundary layer problem with slip, for the semi-infinite flat plate at zero angle of attack may be stated

$$\rho_{\infty} u_{\infty} \frac{\partial u}{\partial x} = \mu \frac{\partial^2 u}{\partial y^2} \quad (34)$$

$$u \rightarrow u_{\infty} \text{ as } y \rightarrow \infty, \quad x > 0 \quad (35)$$

$$u = \zeta \frac{\partial u}{\partial y} \text{ at } y = 0, \quad x > 0 \quad (36)$$

This is reminiscent of the Rayleigh impulsive plate problem, and if condition (35) is replaced by

$$\left. \begin{aligned} u &= u_{\infty} \text{ at } x = 0, \quad y > 0 \\ u &\text{ bounded in the entire plane} \end{aligned} \right\} \quad (37)$$

one may solve this new problem with the aid of Laplace transforms (reference 7). Fortunately, the solution to this second problem satisfies conditions (35) so the solution to the original problem is at hand. This solution is:

$$u = u_{\infty} \left[\operatorname{erfc} \left(\sqrt{\frac{u_{\infty} y^2}{4 \nu_{\infty} x}} \right) + \exp \left(\frac{\nu_{\infty} x}{u_{\infty} \zeta^2} + \frac{y}{\zeta} \right) \operatorname{erfc} \left(\sqrt{\frac{\nu_{\infty} x}{u_{\infty} \zeta^2}} + \sqrt{\frac{u_{\infty} y^2}{4 \nu_{\infty} x}} \right) \right] \quad (38)$$

$$\begin{aligned} \frac{\partial u}{\partial x} &= u_{\infty} \left[\frac{\nu_{\infty}}{u_{\infty} \zeta} \exp \left(\frac{\nu_{\infty} x}{u_{\infty} \zeta^2} + \frac{y}{\zeta} \right) \operatorname{erfc} \left(\sqrt{\frac{\nu_{\infty} x}{u_{\infty} \zeta^2}} + \sqrt{\frac{u_{\infty} y^2}{4 \nu_{\infty} x}} \right) \right. \\ &\quad \left. - \frac{1}{\sqrt{\pi}} \sqrt{\frac{\nu_{\infty}}{u_{\infty} \zeta^2 x}} \exp \left(\frac{-u_{\infty} y^2}{4 \nu_{\infty} x} \right) \right] \end{aligned} \quad (39)$$

$$\frac{\partial u}{\partial y} = u_{\infty} \left[\frac{1}{\zeta} \exp\left(\frac{\nu_{\infty} x}{u_{\infty} \zeta^2} + \frac{y}{\zeta}\right) \operatorname{erfc}\left(\sqrt{\frac{\nu_{\infty} x}{u_{\infty} \zeta^2}} + \sqrt{\frac{u_{\infty} y^2}{4 \nu_{\infty} x}}\right) \right] \quad (40)$$

$$\frac{\partial^2 u}{\partial y^2} = \frac{u_{\infty}}{\nu_{\infty}} \frac{\partial u}{\partial x} \quad (41)$$

If one defines a local skin friction coefficient,

$$C_f = \frac{\mu}{\frac{1}{2} \rho_{\infty} u_{\infty}^2} \left(\frac{\partial u}{\partial y} \right)_{y=0} \quad (42)$$

one obtains, for this problem,

$$C_f = \frac{2 \nu_{\infty}}{u_{\infty} \zeta} \exp\left(\frac{\nu_{\infty} x}{u_{\infty} \zeta^2}\right) \operatorname{erfc}\left(\sqrt{\frac{\nu_{\infty} x}{u_{\infty} \zeta^2}}\right) \quad (43)$$

The drag coefficient for that portion of the plate extending from the origin to a distance L downstream will be defined (including both upper and lower plate surfaces) as

$$C_D = \frac{2}{L} \int_0^L C_f dx \quad (44)$$

and for this problem

$$C_D = 4 \sqrt{\frac{\nu_{\infty}}{u_{\infty} L}} \left\{ \sqrt{\frac{u_{\infty} \zeta^2}{\nu_{\infty} L}} \left[\exp\left(\frac{\nu_{\infty} L}{u_{\infty} \zeta^2}\right) \operatorname{erfc}\left(\sqrt{\frac{\nu_{\infty} L}{u_{\infty} \zeta^2}}\right) - 1 \right] + \frac{2}{\sqrt{\pi}} \right\} \quad (45)$$

Equations (38), (43), (45) constitute the solution of the originally posed problem.

An attempt to solve this same problem by a formal iteration in ζ , commences with the formal series

$$u(x, y, \zeta) = u_0(x, y) + \zeta u_1(x, y) + \zeta^2 u_2(x, y) + \dots \quad (46)$$

This yields the iteration scheme

$$\left. \begin{aligned} \rho_{\infty} u_{\infty} \frac{\partial u_0}{\partial x} &= \mu \frac{\partial^2 u_0}{\partial y^2} \\ u_0 &\rightarrow u_{\infty} \text{ as } y \rightarrow \infty, x > 0 \\ u_0 &= 0 \text{ at } y = 0, x > 0 \end{aligned} \right\} \quad (47)$$

and for $n = 1, 2, 3, \dots$

$$\left. \begin{aligned} \rho_{\infty} u_{\infty} \frac{\partial u_n}{\partial x} &= \mu \frac{\partial^2 u_n}{\partial y^2} \\ u_n &\rightarrow 0 \text{ as } y \rightarrow \infty, x > 0 \\ u_n &= \frac{\partial u_{n-1}}{\partial y} \text{ at } y = 0, x > 0 \end{aligned} \right\} \quad (48)$$

The technique employed in solving the problem delineated by equation (34), under the conditions (35) and (36) may be employed here to solve equations (47). The no-slip solution, $u_0(x, y)$ is

$$u_0(x, y) = u_{\infty} \operatorname{erf} \left(\sqrt{\frac{u_{\infty} y^2}{4 \nu_{\infty} x}} \right) \quad (49)$$

It is easy to see that

$$u_n(x, y) = \frac{\partial u_{n-1}(x, y)}{\partial y} \quad \text{for } n = 1, 2, 3, \dots \quad (50)$$

solves equations (48). Thus, formally,

$$u(x, y, \tau) = \sum_{n=0}^{\infty} \tau^n \frac{\partial^n}{\partial y^n} \left(u_{\infty} \operatorname{erf} \left(\sqrt{\frac{u_{\infty} y^2}{4 \nu_{\infty} x}} \right) \right) \quad (51)$$

From the definition (42) and equation (51) one obtains, formally

$$\begin{aligned} C_f &= \frac{\mu}{\frac{1}{2} \rho_{\infty} u_{\infty}^2} \sum_{n=0}^{\infty} \tau^n \left[\frac{\partial^{n+1}}{\partial y^{n+1}} \left(u_{\infty} \operatorname{erf} \left(\sqrt{\frac{u_{\infty} y^2}{4 \nu_{\infty} x}} \right) \right) \right]_{y=0} \\ &= \frac{2}{\sqrt{\pi}} \sqrt{\frac{\nu_{\infty}}{u_{\infty} x}} \sum_{n=0}^{\infty} \tau^n \left[\frac{\partial^n}{\partial y^n} \exp \left(\frac{-u_{\infty} y^2}{4 \nu_{\infty} x} \right) \right]_{y=0} \end{aligned} \quad (52)$$

$$= \frac{2}{\pi} \sqrt{\frac{\nu_{\infty}}{u_{\infty} x}} \sum_{n=0}^{\infty} (-1)^n \Gamma\left(\frac{2n+1}{2}\right) \left(\frac{u_{\infty} \zeta^2}{\nu_{\infty} x}\right)^n$$

Integration by parts and the use of induction on expression (43) yields

$$C_f \sim \frac{2}{\pi} \sqrt{\frac{\nu_{\infty}}{u_{\infty} x}} \sum_{n=0}^{\infty} (-1)^n \Gamma\left(\frac{2n+1}{2}\right) \left(\frac{u_{\infty} \zeta^2}{\nu_{\infty} x}\right)^n \quad (53)$$

Thus, one sees that the formal iteration scheme led to an asymptotic (around $\zeta = 0$) expansion for C_f (which is uniform in x for x bounded away from zero). Expression (52) is useless at $x = 0$ and cannot help in the determination of C_D ; it can, however, in a series of unjustified steps, be summed exponentially, to yield expression (43) (see Appendix III).

From expression (43), one learns that

$$\left. \begin{aligned} \lim_{x \rightarrow 0} C_f &= \frac{2 \nu_{\infty}}{u_{\infty} \zeta} && \text{for fixed } \zeta > 0 \\ \lim_{\zeta \rightarrow 0} C_f &= \frac{2}{\sqrt{\pi}} \sqrt{\frac{\nu_{\infty}}{u_{\infty} x}} && \text{for fixed } x > 0 \end{aligned} \right\} \quad (54)$$

and the non-uniformity of C_f in ζ and x at $\zeta = x = 0$ is apparent.

From expression (53),

$$C_f \sim \frac{2}{\sqrt{\pi}} \sqrt{\frac{\nu_{\infty}}{u_{\infty} x}} \left[1 - \frac{1}{2} \frac{u_{\infty} \zeta^2}{\nu_{\infty} x} + \frac{3}{4} \left(\frac{u_{\infty} \zeta^2}{\nu_{\infty} x} \right)^2 + \dots \right] \quad (55)$$

Integration by parts and the use of induction on expression (45) yields

$$\begin{aligned} C_D \sim \frac{8}{\sqrt{\pi}} \sqrt{\frac{\nu_{\infty}}{u_{\infty} L}} &\left[1 - \frac{\sqrt{\pi}}{2} \sqrt{\frac{u_{\infty} \zeta^2}{\nu_{\infty} L}} + \frac{1}{2} \frac{u_{\infty} \zeta^2}{\nu_{\infty} L} \right. \\ &\left. - \frac{1}{4} \left(\frac{u_{\infty} \zeta^2}{\nu_{\infty} L} \right)^2 + \dots \right] \end{aligned} \quad (56)$$

From equations (55) and (56) it is evident that a slip correction to the local skin friction of order ζ^2 does not imply that the slip correction to the drag is also of order ζ^2 , since the non-uniformity is sufficiently strong so that the actual correction is of order ζ .

3.0 CONCLUSIONS

1. The scheme, which Alden used to correct the Blasius solution, cannot lead to a solution, even formally, of the incompressible Navier-Stokes semi-infinite flat plate problem.
2. For the incompressible linearized flat plate problem the inclusion of a slip boundary condition introduces a correction to the local skin friction of order γ^2 but, due to a non-uniformity in the solution, corrects the drag coefficient to order γ .

APPENDIX I

$$F_2(\xi) = 4\xi f_1''' - \xi^2 f_1'''' - \xi^2 f_1 f_1''' - 2\xi f_1 f_1'' + \xi^2 f_1' f_1''$$

$$\begin{aligned} F_3(\xi) = & -3f_2' f_2'' + f_2 f_2''' - 10f_1' f_2 - \xi^2 f_1' f_1'' - 2f_1 f_2' - \xi^2 f_1 f_2''' \\ & - 2\xi f_1'' f_2 + \xi^2 f_2' f_1'' + \xi^2 f_2 f_1''' + 2\xi f_2'' f_1 - 12 f_2'' \\ & + 4\xi f_2''' - \xi^2 f_2'''' \end{aligned}$$

$$\begin{aligned} F_4(\xi) = & -3f_2' f_3'' + f_2 f_3''' - 2f_2'' f_3' - f_3' f_2'' + 3f_3 f_2''' + 2f_3'' f_2' \\ & - 60f_1 f_3 - 3\xi^2 f_1' f_3'' - 12f_1 f_3' - \xi^2 f_1 f_3''' - 6\xi f_1'' f_3 \\ & - 6f_2' f_2 - \xi^2 f_2' f_2'' + 2f_2 f_2' + \xi^2 f_2 f_2''' - 4f_2 f_2' - 2\xi f_2'' f_2' \\ & + \xi^2 f_3' f_1'' + 3\xi^2 f_3 f_1''' - 24f_3 f_1' + 2\xi f_3'' f_1 - 6\xi^2 f_1' f_2 \\ & - 2\xi^2 f_1 f_2' + 4\xi f_2 f_1 - 40f_3'' + 4\xi f_3''' - \xi^2 f_3'''' - 56f_2 \\ & + 8\xi f_2' - 4\xi^2 f_2'' \end{aligned}$$

APPENDIX II

From Appendix I and equation (21) $F_2 \sim 0$. Introducing the integration constants, K_{mn} , one obtains

$$F_2 \sim 0$$

$$F_{21} \sim C_{21} + K_{21}$$

$$F_{22} \sim (C_{21} + K_{21})(\xi - \beta/2) + C_{22} + K_{22}$$

$$h_2 = 1$$

$$F_{23} \sim (C_{21} + K_{21}) \frac{(\xi - \beta/2)^2}{2} + (C_{22} + K_{22})(\xi - \beta/2) + K_{23} \quad (\text{II-1})$$

$$F_{24} \sim F_{23}$$

$$f_2 \sim \frac{C_{21} + K_{21}}{2} \left[\frac{(\xi - \beta/2)}{2} + O(\xi^{-1}) \right] + \frac{C_{22} + K_{22}}{2} + \frac{K_{23}}{2} \left[(\xi - \beta/2)^{-1} + \frac{(\xi - \beta/2)^{-3}}{2} \right] + O(\xi^{-5})$$

Because of conditions (16), one must put

$$C_{21} = -K_{21} \quad (\text{II-2})$$

Using equations (14), (24), (25) and (28), one obtains

$$f_2 = \frac{C_{22} + K_{22}}{2} + K_{23} \left[\frac{(\xi - \beta/2)^{-1}}{2} + \frac{(\xi - \beta/2)^{-3}}{4} \right] + O(\xi^{-5})$$

$$f_2' = -K_{23} \left[\frac{(\xi - \beta/2)^{-2}}{2} + \frac{3(\xi - \beta/2)^{-4}}{4} \right] + O(\xi^{-6})$$

$$f_2'' = K_{23} \left[(\xi - \beta/2)^{-3} + 3(\xi - \beta/2)^{-5} \right] + O(\xi^{-7}) \quad (\text{II-3})$$

$$f_2''' = -K_{23} \left[3(\xi - \beta/2)^{-4} + 15(\xi - \beta/2)^{-6} \right] + O(\xi^{-8})$$

$$f_2^{(4)} = K_{23} \left[12(\xi - \beta/2)^{-5} + 90(\xi - \beta/2)^{-7} \right] + O(\xi^{-9})$$

Appendix I and equations (21) and (II-3) yield

$$F_3 = -10(C_{22} + K_{22}) + 6\beta K_{23}(\xi - \beta/2)^{-2} + O(\xi^{-3}) \quad (\text{II-4})$$

Proceeding in the same way as with F_2 ,

$$F_{31} = -10(C_{22} + K_{22})(\xi - \beta/2) + C_{31} + K_{31} \\ - 6\beta K_{23}(\xi - \beta/2)^{-1} + O(\xi^{-2})$$

$$F_{32} = -5(C_{22} + K_{22})(\xi - \beta/2)^2 + (C_{31} + K_{31})(\xi - \beta/2) \\ - 6\beta K_{23} \ln(\xi - \beta/2) + O(\xi^{-1})$$

$$h_3 = 4(\xi - \beta/2)^2 - 2 \quad (\text{II-5})$$

$$F_{33} = -4(C_{22} + K_{22})(\xi - \beta/2)^5 [1 + O(\xi^{-2})] \\ + (C_{31} + K_{31})(\xi - \beta/2)^4 [1 + O(\xi^{-2})] \\ - 8\beta K_{23}(\xi - \beta/2)^3 \ln(\xi - \beta/2) [1 + O(\xi^{-2})] + O(\xi^3)$$

$$[h_3(\xi)]^{-2} = \frac{1}{16}(\xi - \beta/2)^{-4} + O(\xi^{-6})$$

$$\begin{aligned}
F_{34} = & -\frac{1}{4}(C_{22}+K_{22})(\xi-\beta/2)\left[1+O(\xi^{-2})\right] + \frac{(C_{31}+K_{31})}{16}\left[1+O(\xi^{-2})\right] \\
& -\frac{1}{2}\beta K_{23}(\xi-\beta/2)^{-1}\ln(\xi-\beta/2)\left[1+O(\xi^{-2})\right] \\
& + O(\xi^{-1})
\end{aligned}$$

$$\begin{aligned}
f_3 = & -\frac{1}{2}(C_{22}+K_{22})(\xi-\beta/2)^2\left[1+O(\xi^{-2})\right] \\
& + \frac{(C_{31}+K_{31})}{8}(\xi-\beta/2)\left[1+O(\xi^{-2})\right] \\
& -\beta K_{23}\ln(\xi-\beta/2)\left[1+O(\xi^{-2})\right] \\
& + O(1)
\end{aligned} \tag{II-6}$$

Because of conditions (16),

$$\begin{aligned}
C_{22} &= -K_{22} \\
C_{31} &= -K_{31}
\end{aligned} \tag{II-7}$$

$$\begin{aligned}
f_3 &= -\beta K_{23}\ln(\xi-\beta/2) + O(1) + \ln(\xi-\beta/2)O(\xi^{-2}) \\
f_3' &= -\beta K_{23}(\xi-\beta/2)^{-1} + O(\xi^{-2}) + \ln(\xi-\beta/2)O(\xi^{-3}) \\
f_3'' &= \beta K_{23}(\xi-\beta/2)^{-2} + O(\xi^{-3}) + \ln(\xi-\beta/2)O(\xi^{-4}) \\
f_3''' &= -2\beta K_{23}(\xi-\beta/2)^{-3} + O(\xi^{-4}) + \ln(\xi-\beta/2)O(\xi^{-5}) \\
f_3^{(4)} &= 6\beta K_{23}(\xi-\beta/2)^{-4} + O(\xi^{-5}) + \ln(\xi-\beta/2)O(\xi^{-6})
\end{aligned} \tag{II-8}$$

$$F_4 = 168\beta K_{23} \ln(\xi - \beta/2) + O(1) + \ln(\xi - \beta/2) O(\xi^{-2})$$

$$F_{41} = 168\beta K_{23} (\xi - \beta/2) \ln(\xi - \beta/2) + O(\xi) + \ln(\xi - \beta/2) O(1)$$

$$F_{42} = 84\beta K_{23} (\xi - \beta/2)^2 \ln(\xi - \beta/2) + [\ln(\xi - \beta/2)]^2 O(1) \\ + O(\xi^2) + \ln(\xi - \beta/2) O(\xi)$$

$$h_4 = 16(\xi - \beta/2)^4 - 48(\xi - \beta/2)^2 + 12$$

$$F_{43} = 192\beta K_{23} (\xi - \beta/2)^7 \ln(\xi - \beta/2) \\ + [\ln(\xi - \beta/2)]^2 O(\xi^5) + O(\xi^7) \\ + \ln(\xi - \beta/2) O(\xi^6)$$

$$F_{44} = \frac{3}{4}\beta K_{23} (\xi - \beta/2)^8 \ln(\xi - \beta/2) \\ + [\ln(\xi - \beta/2)]^2 O(\xi^{-3}) + O(\xi^{-1}) \\ + \ln(\xi - \beta/2) O(\xi^{-2})$$

(II-9)

$$f_4 = 6\beta K_{23} (\xi - \beta/2)^2 \ln(\xi - \beta/2) \\ + [\ln(\xi - \beta/2)]^2 O(1) + O(\xi^2) \\ + \ln(\xi - \beta/2) O(\xi)$$

$$[h_4(\xi)]^{-2} = \frac{1}{256} (\xi - \beta/2)^{-8} + O(\xi^{-10})$$

APPENDIX III

Proceeding from equation (52) in a purely formal manner:

$$\begin{aligned}
 C_f &= \frac{2}{\pi} \sqrt{\frac{\nu_{\infty}}{u_{\infty} x}} \sum_{n=0}^{\infty} \Gamma\left(\frac{2n+1}{2}\right) \left(\frac{-u_{\infty} \zeta^2}{\nu_{\infty} x}\right)^n \left[\frac{1}{\Gamma(n+1)} \int_0^{\infty} e^{-t} t^n dt \right] \\
 &= \frac{2}{\pi} \sqrt{\frac{\nu_{\infty}}{u_{\infty} x}} \int_0^{\infty} e^{-t} \left[\sum_{n=0}^{\infty} \frac{\Gamma\left(\frac{2n+1}{2}\right)}{\Gamma(n+1)} \left(\frac{-u_{\infty} \zeta^2 t}{\nu_{\infty} x}\right)^n \right] dt \\
 &= \frac{2}{\pi} \frac{\nu_{\infty}}{u_{\infty} \zeta} \sqrt{\frac{u_{\infty} \zeta^2}{\nu_{\infty} x}} \int_0^{\infty} e^{-t} \left[1 + \frac{u_{\infty} \zeta^2}{\nu_{\infty} x} t \right]^{-\frac{1}{2}} dt
 \end{aligned} \tag{III-1}$$

The expression

$$I(\lambda) = \frac{1}{\lambda} \int_0^{\infty} e^{-t} \left[1 + \frac{t}{\lambda^2} \right]^{-\frac{1}{2}} dt \tag{III-2}$$

with

$$\lambda = \sqrt{\frac{u_{\infty} x}{u_{\infty} \zeta^2}} \tag{III-3}$$

satisfies

$$\frac{dI}{d\lambda} - 2\lambda I + 2 = 0 \tag{III-4}$$

$$I(0) = \sqrt{\pi}$$

Therefore

$$I(\lambda) = \sqrt{\pi} \exp(\lambda^2) \operatorname{erfc}(\lambda) \tag{III-5}$$

and finally,

$$C_f = \frac{2V_{\infty}}{u_{\infty} \zeta} \exp\left(\frac{V_{\infty} x}{u_{\infty} \zeta^2}\right) \operatorname{erfc}\left(\sqrt{\frac{V_{\infty} x}{u_{\infty} \zeta^2}}\right) \quad (\text{III-6})$$

This method of "exponentially summing" a divergent series is discussed by Borel in reference 16.

REFERENCES

1. Goldstein, S., "Modern Developments in Fluid Dynamics", Oxford, 1938, pp. 135-138.
2. Blasius, H., Zeitschrift fur Math. u. Physik. Vol. 56, 1908, pp. 4-13.
3. von Karman, Th., Bulletin of the American Mathematical Society, 1940, pp. 655-659.
4. Alden, H. L., "Second Approximation to the Laminar Boundary Layer Flow Over a Flat Plate", Jour. Math. and Phys., April 1948, p. 91.
5. Lin, T. C., and Schaaf, S. A. "The Effect of Slip on Flow Near a Stagnation Point and in a Boundary Layer", NACA TN No. 2568, December, 1951.
6. Maslen, S. A., "Second Approximation to Laminar Boundary Layer Flow on Flat Plate in Slip Flow", NACA TN No. 2818, November 1952.
7. Schaaf, S. A., and Sherman, F. S., "Skin Friction in Slip Flow", Jour. of the Aeronautical Sciences, Vol. 21, No. 2, February, 1954.
8. Schamberg, R., "The Fundamental Differential Equations and the Boundary Conditions for High Speed Slip-Flow, and their Application to Several Specific Problems", C. I. T. Thesis, 1947.
9. Kennard, E. H., "Kinetic Theory of Gases", First ed., McGraw-Hill Book Co., Inc., 1938, p. 296.
10. Lamb, H., "Hydrodynamics", Dover, 1945, pp. 684-686.
11. Kaplun, Saul, "The Role of Coordinate Systems in Boundary-Layer Theory", Z. fur Angew. Math. u. Phys., Vol. 5 1954, p. 111.
12. von Mises, R. and Friedrichs, K. O., "Fluid Dynamics", Brown University Notes, 1941.
13. Latta, G. E., "An Investigation of the Classical Boundary Layer Associated with Oseen's Equation for the Flow of a Viscous Incompressible Fluid Past a Semi-Infinite Flat Plate at Zero Angle of Attack", Preliminary write-up, not yet published.
14. Weyl, Hermann, "On the Differential Equations of the Simplest Boundary-Layer Problems", Ann. of Math., Vol. 43, 1942, p. 381.
15. Prandtl, L., "Aerodynamic Theory", (Edited by W. F. Durand), Vol. 3, pp. 84-90.
16. Borel, E. "Lecons sur les Series Divergentes", Paris, 1901, p. 97.

DISTRIBUTION LIST FOR ONR TECHNICAL REPORTS

Chief of Naval Research Department of the Navy Washington 25, D. C. Attn: Code 438	(2)	Chief, Bureau of Ordnance Department of the Navy Washington 25, D. C. Attn: Research and Development Division	(1)
Commanding Officer Office of Naval Research Branch Office 495 Summer Street Boston, Massachusetts	(0)	Office of Ordnance Research Department of the Army Washington 25, D. C.	(1)
Commanding Officer Office of Naval Research Branch Office The John Crerar Library Bldg. 86 East Randolph Street Chicago 1, Illinois	(1)	Commander Air Research and Development Command Office of Scientific Research P. O. Box 1395 Baltimore 18, Maryland Attn: Fluid Mech. Div.	(1)
Commanding Officer Office of Naval Research Branch Office 346 Broadway New York 13, N. Y.	(1)	Director of Research National Advisory Committee for Aeronautics 1512 H Street, N. W. Washington 25, D. C.	(1)
Commanding Officer Office of Naval Research Branch Office 1030 East Green Street Pasadena 1, California	(1)	Director Langley Aeronautical Laboratory National Advisory Committee for Aeronautics Langley Field, Virginia	(1)
Commanding Officer Office of Naval Research Branch Office 1000 Geary Street San Francisco 24, California	(1)	Director National Bureau of Standards Washington 25, D. C. Attn: Fluid Mechanics Section	(1)
Commanding Officer Office of Naval Research Navy #100, Fleet Post Office New York, N. Y.	(3)	Professor R. Courant Institute for Math. and Mechanics New York University 45 Fourth Avenue New York 3, N. Y.	(1)
Director Naval Research Laboratory Washington 25, D. C. Attn: Code 2021	(6)	Professor G. Kuerti Department of Mechanical Eng'g Case Institute of Technology Cleveland, Ohio	(1)
Documents Service Center Armed Services Technical Information Agency Knott Building Dayton 2, Ohio	(5)	Professor W. R. Sears, Director Graduate School of Aero. Eng'g Cornell University Ithaca, New York	(1)
Chief, Bureau of Aeronautics Department of the Navy Washington 25, D. C. Attn: Research Division	(1)	Commanding Officer Naval Ordnance Laboratory White Oak, Maryland Attn: Aeroballistics Res. Dept.	(1)

Commanding Officer and Director
David Taylor Model Basin
Washington 7, D. C.
Attn: Aerodynamics Laboratory (1)

Ballistics Research Laboratory
Department of the Army
Aberdeen Proving Ground
Aberdeen, Maryland
Attn: Mr. R. H. Kent (1)

Professor F. H. Clauser
Department of Aeronautics
Johns Hopkins University
Baltimore 18, Maryland (1)

Professor J. R. Markham
Department of Aeronautical Eng'g
Massachusetts Institute of Technology
Cambridge 39, Massachusetts (1)

Professor J. F. Ludloff
Guggenheim School of Aeronautics
New York University
New York 53, N. Y. (1)

Professor A. Kantrowitz
Graduate School of Aero. Eng'g
Cornell University
Ithaca, New York (1)

Bureau of Aeronautics General
Representative USN
Central District
Wright-Patterson Air Force Base
Dayton, Ohio
Attn: WCOWS-3 (Mr. James G. Law)(1)

Commanding General
Air Force Cambridge Research Center
230 Albany Street
Cambridge 39, Massachusetts
Attn: Dr. H. D. Edwards
Geophysics Division (1)

Dr. Morton Alperin
Chief, Western Regional Office
Air Research and Development Center
P. O. Box 2035
Pasadena 2, California (1)

Chief
Armed Forces Special Weapons Project
P. O. Box 2610
Washington 25, D. C. (1)

Commanding Officer
Arnold Engineering Development Center
Tullahoma, Tennessee
Attn: ACGT, ACR, AEMCL (3)

Mr. C. K. Bauer
Hermes Project
General Electric Company
Schenectady 5, New York (1)

Professor W. S. Bradfield
Department of Aeronautical Eng'g
University of Minnesota
Minneapolis, Minnesota (1)

Chief, Bureau of Aeronautics
Department of the Navy
Washington 25, D. C.
Attn: Aero. and Hydro. Branch (1)

Chief of Staff
Hdqs, USAF
Directorate of Research and Develop.
Washington, D. C.
Attn: AFDRD-AC2 (1)

Professor R. E. Street
University of Washington
Department of Aeronautical Eng'g
Seattle 5, Washington (1)

Chief of Naval Research
Department of the Navy
Washington 25, D. C.
Attn: Code 419 (Dr. I. Estermann)(1)

Chief, Bureau of Ordnance
Department of the Navy
Washington 25, D. C.
Attn: Code Re2c (1)

Commanding Officer
Naval Ordnance Laboratory
White Oak, Maryland
Attn: Hyperballistics Div. (1)

Ballistics Research Laboratory
Department of the Army
Aberdeen Proving Ground
Aberdeen, Maryland
Attn: Mr. C. W. Lampson (1)

Exterior Ballistics Laboratory
Department of the Army
Aberdeen Proving Ground
Aberdeen, Maryland
Attn: Mr. F. D. Bennett (1)

Professor J. Kestin
Division of Engineering
Brown University
Providence 12, Rhode Island (1)

Graduate Division of
Applied Mathematics
Brown University
Providence 12, Rhode Island (1)

Professor D. F. Hornig
Metcalf Research Laboratory
Brown University
Providence 12, Rhode Island (1)

Professor Hsue-Shien Tsien
Guggenheim Jet Propulsion Center
California Institute of Technology
Pasadena 4, California (1)

Professor A. B. Arons
Department of Physics
Amherst College
Amherst, Massachusetts (1)

Professor Walker Bleakney
Palmer Physical Laboratory
Princeton University
Princeton, New Jersey (1)

Professor George F. Carrier
Department of Engineering Sciences
Harvard University
Pierce Hall
Cambridge 38, Massachusetts (1)

Professor N. S. Plesset
Hydrodynamics Laboratory
California Institute of Technology
Pasadena 4, California (1)

Professor Howard W. Emmons
Department of Engineering Sciences
Harvard University
Pierce Hall
Cambridge 38, Massachusetts (1)

Dr. R. N. Hollyer, Jr.
Applied Physics Laboratory
Johns Hopkins University
8621 Georgia Avenue
Silver Spring, Maryland (1)

Professor H. G. Stever
Department of Aeronautical Eng'g
Massachusetts Institute of Technology
Cambridge 39, Massachusetts (1)

Professor R. G. Stoner
Department of Physics
Pennsylvania State University
State College, Pennsylvania (1)

Professor S. F. Borg
Department of Civil Engineering
Stevens Institute of Technology
711 Hudson Street
Hoboken, New Jersey (1)

Professor E. L. Resler
Institute for Fluid Dynamics
and Applied Mathematics
University of Maryland
College Park, Maryland (1)

Professor Otto Laporte
Engineering Research Institute
East Engineering Building
University of Michigan
Ann Arbor, Michigan (1)

Professor J. G. Kirkwood
Department of Chemistry
Yale University
New Haven, Connecticut (1)

Mr. K. L. Sandefur
Department of Physics
Midwest Research Institute
4049 Pennsylvania Avenue
Kansas City, Missouri (1)

Dr. S. J. Fraenkel
Armour Research Institute
Illinois Institute of Technology
Chicago 16, Illinois (1)

Chief, Bureau of Aeronautics
Department of the Navy
Washington 25, D. C.
Attn: Airframe Design Div. (1)

Chief, Bureau of Ordnance
Department of the Navy
Washington 25, D. C.
Attn: Code Re9a (1)

Commanding Officer Naval Ordnance Laboratory White Oak, Maryland Attn: Aerophysics Division (1)	Professor C. C. Lin Department of Mathematics Massachusetts Institute of Technology Cambridge 39, Massachusetts (1)
Commander Naval Ordnance Test Station Inyokern China Lake, California (1)	Massachusetts Institute of Technology Department of Mechanical Engineering Cambridge 39, Massachusetts Attn: Professor J. Kaye Professor A. H. Shapiro Professor E. S. Taylor (3)
Director Ames Aeronautical Laboratory National Advisory Committee for Aeronautics Moffett Field, California (1)	Professor H. G. Lew Department of Aeronautical Engineering Pennsylvania State University State College, Pennsylvania (1)
Director Lewis Flight Propulsion Laboratory National Advisory Committee for Aeronautics 21000 Brookpark Road Cleveland 11, Ohio (1)	Stanford University Guggenheim Aeronautical Laboratory Stanford, California (1)
Professor A. Ferri Department of Aeronautical Engineering and Applied Mechanics Polytechnic Institute of Brooklyn 99 Livingston Street Brooklyn 2, New York (1)	Professor S. A. Schaaf Low Pressures Research Project University of California Berkeley, California (1)
Brown University Graduate Division of Applied Math. Providence 12, Rhode Island (1)	Dean L. M. K. Boelter University of California Los Angeles 24, California (1)
California Institute of Technology Guggenheim Aeronautical Laboratory Pasadena 4, California Attn: Professor P. A. Lagerstrom Professor L. Lees Professor H. W. Liepmann (3)	Institute for Fluid Dynamics and Applied Mathematics University of Maryland College Park, Maryland (1)
Professor S. Corrsin Department of Aeronautical Engineering Johns Hopkins University Baltimore 18, Maryland (1)	Professor A. M. Kueth Department of Aeronautical Eng'g University of Michigan Ann Arbor, Michigan (1)
Professor P. F. Meader Division of Engineering Brown University Providence 12, Rhode Island (1)	Professor J. D. Akerman Department of Aeronautical Eng'g University of Minnesota Minneapolis, Minnesota (1)
Dr. F. N. Frenkiel Applied Physics Laboratory Johns Hopkins University 8621 Georgia Avenue Silver Spring, Maryland (1)	Professor C. H. Fletcher Department of Aeronautical Eng'g University of Illinois Urbana, Illinois (1)
	Professor M. J. Thompson Defense Research Laboratory University of Texas Austin, Texas (1)

Director
Jet Propulsion Laboratory
California Institute of Technology
Pasadena 4, California (1)

Professor R. P. Harrington
Department of Aeronautical Eng'g
Rensselaer Polytechnic Institute
Troy, New York (1)

Professor J. V. Charyk
Forrestal Research Center
Princeton University
Princeton, New Jersey (1)

Professor S. M. Bogdonoff
Department of Aeronautical Eng'g
Princeton University
Princeton, New Jersey (1)

Dr. Ronald F. Probst
Division of Engineering
Brown University
Providence, Rhode Island (1)

Professor M. U. Clauser
Department of Aeronautical Eng'g
Purdue University
Lafayette, Indiana (1)

Dr. J. M. Jones
Engineering Research Institute
University of Michigan
Ann Arbor, Michigan (1)

Dr. J. Kaplan
Department of Physics
University of California
Los Angeles, California (1)

Professor J. H. Keenan
Mechanical Engineering Department
Massachusetts Institute of Technology
Cambridge 39, Massachusetts (1)

Professor M. Z. Kryzwoblocki
University of Illinois
Urbana, Illinois (1)

Dr. A. R. Kuhltau
Department of Physics
University of Virginia
Charlottesville, Virginia (1)

Professor H. Lewy
Applied Math Group, Math. Department
University of California
Berkeley, California (1)

Dr. James E. Lipp
The Rand Corporation
1500 Fourth Street
Santa Monica, California (1)

Dr. Henry T. Nagamatsu, Director
Hypersonic Wind Tunnel
California Institute of Technology
Pasadena 4, California (1)

Radiation Laboratory
University of California
Berkeley 4, California
Attn: R. Wakerling (1)

Mr. R. H. Shick
Consolidated Vultee Aircraft Corp.
San Diego 12, California (1)

Sverdrup and Parcel, Inc.
Consulting Engineers
Syndicate Trust Building
St. Louis, Missouri (1)

Dr. Theodore Theodorsen
University of Maryland
Building J, Room 314
College Park, Maryland (1)

Western Coordination Office
National Advisory Committee
for Aeronautics
7660 Beverly Blvd.
Los Angeles 36, California (1)

Dr. E. P. Williams
Missiles Division
Rand Corporation
1500 Fourth Street
Santa Monica, California (1)

Commanding General
Wright Air Development Center
Wright-Patterson Air Force Base
Dayton, Ohio

Attn: WCRRS-3 (1)
MCIDXE (1)

Commander
Naval Ordnance Test Station
Inyokern
China Lake, California
Attn: Technical Library (1)

Commander
Wright Air Development Center
Wright-Patterson Air Force Base
Dayton, Ohio
Attn: WCKSC (0)

Glenn L. Martin Company
Baltimore 3, Maryland
Attn: C. J. Pierce (1)

Professor R. G. Folsom
Engineering Research Institute
University of Michigan
Ann Arbor, Michigan (1)

J. R. Stalder
1831 Hamilton Avenue
Palo Alto, California (1)

Professor Paul A. Libby
Dept. of Aero. Engineering
Polytechnic Institute of Brooklyn
99 Livingston Street
Brooklyn, New York (1)

Professor K. Stewartson
Department of Mathematics
Bristol University
Bristol, England (1)

Professor Harold Grad
Institute for Math and Mech.
New York University
45 Fourth Avenue
New York 3, New York

Dr. R. J. Emrich
Department of Physics
Lehigh University
Bethlehem, Pennsylvania (1)

Dr. J. O. Herschfelder
P. O. Box 2127
Madison 6, Wisconsin (1)

Commander
Western Development Division
Post Office Box 262
Inglewood, California (1)